

Question	Answer	Marks	Guidance
1(a)	high melting point/mp/mpt OR high boiling point/bp/bpt; poor/non conductor (when liquid and/or solid);	3	I mpt/bpt above room temp
(b)	(good) conductor when <u>solid</u> (and liquid);	2	A (good) conductor in any state/both states I high melting point/boiling point R low melting point/boiling point
(c)	melting point/ $-7(^{\circ}\text{C})$ is below room temperature/ $25(^{\circ}\text{C})$ /RTP ora; boiling point/ $59(^{\circ}\text{C})$ is above room temperature/ $25(^{\circ}\text{C})$ /RTP ora;	3	I low melting point/boiling point/conductivity 25($^{\circ}\text{C}$)/room temperature/RTP is in between $-7(^{\circ}\text{C})$ and $59(^{\circ}\text{C})$ OR 25($^{\circ}\text{C}$)/room temperature/RTP is between mpt and bpt would both score the 2 evidence marks
(d)	high melting point/mp/mpt OR high boiling point/bp/bpt; BOTH poor/non conductor when solid and good conductor when liquid OR molten/only conduct when liquid;	3	A melting point and boiling point both above room temp/ 25°C /RTP I conducts when aqueous or in solution I conducts in liquid due to free electrons

Question	Answer	Marks	Guidance
2(a)(i)	3;	1	
1(a)(ii)	2O_3 ;	1	As_2O_5
1(a)(iii)	4;	1	
(b)(i)	3 ;	1	
1(b)(ii)	$^{2+}$;	1	
(b)(iii)	$^+$;	1	
(c)	M1 2 double bonds, one between each O and the C atom; M2 each O has 8 outer electrons; M3 each C has 8 outer electrons;	3	R wrong symbols for O for M2 R wrong symbols for C for M3 I missing symbols A any combination of x and o

- 3 (a) (i) $6\text{Li} + \text{N}_2 = 2\text{Li}_3\text{N}$
species (1) balancing (1)
- (ii) N^{3-} ion drawn correctly [1]
Charges correct (minimum 1 \times Li ion and 1 nitride ion) [1]
- (b) 3 \times shared pairs between N and 3 \times F [1]
only 2 non-bonding electrons on N, 6 non-bonding electrons on each F
(COND on first point) [1]
- (ii) Strong attractive forces/strong ionic bonds in lithium nitride [1]
weak (attractive) forces between molecules in NF_3 [1]

[Total: 8]

- 4 (a) (i) incomplete combustion **or** limited oxygen/less oxygen/not enough oxygen (1) [1]
- (ii) any **two** from:
 (forward) reaction is endothermic (1)
 high temperature increases yield/favours forward reaction/shifts equilibrium to right (1)
 faster reaction (rate) (1) [2]
- (iii) any **two** from:
 high pressure reduces yield **or** favours LHS (1)
 because LHS has smaller volume **or** number of moles/number of molecules (of gas) ORA (1)
 (high pressure plant is) expensive/dangerous/explosion/leaks [2]
- 5 (b) hydrogen **and** chlorine/H₂ **and** Cl₂ (1)
 sodium hydroxide/NaOH/Na⁺OH (1)
 $2\text{H}^+ + 2\text{e} \rightarrow \text{H}_2/2\text{H}^+ \rightarrow \text{H}_2 - 2\text{e}$ (1)
 $2\text{Cl} \rightarrow \text{Cl}_2 + 2\text{e}/2\text{Cl} - 2\text{e} \rightarrow \text{Cl}_2$ (1)
 Hydrogen/H₂/H/H⁺ at cathode **and** chlorine/chloride/Cl₂/Cl/Cl at anode (1) [5]
- 5 (c) each chlorine 1 bond pair and 3 non-bond pair (1)
 oxygen atom 2 non-bond pairs and 2 bond pairs as double bond (1)
 carbon atom 4 bond pairs including 2 bond pairs as double bond (1) [3]

[Total: 13]

- 5 (a) (i) any ambiguous formula, e.g. $\text{GeH}_3\text{-GeH}_2\text{-GeH}_3$ [1]
- (ii) $\text{Ge}_n\text{H}_{2n+2}$ [1]
NOT C instead of Ge
- (b) correct formula
COND 4bps around germanium atom [1]
COND 3nbps and 1bp around each chlorine atom [1]
- (c) four oxygen atoms around each germanium atom [1]
two germanium atoms around each oxygen atom [1]
tetrahedral [1]
- (d) oxidation [1]
COND increase in oxidation number [1]
ACCEPT: electron loss